



CHLORINE SANITIZER

For Automatic Dish Machines

This product is a chlorine bearing, sodium hypochlorite solution for sanitization in the Dairy, Food Processing and Food Service industries. It is an effective sanitizer for utensils, glassware and as a terminal sanitizing rinse in machine dishwashing. This product may also be used for fruit and vegetable washing.

■ PHYSICAL PROPERTIES

APPEARANCE: Light Yellow Liquid

SPECIFIC GRAVITY: 1.023

ODOR: Bleach

pH (Conc.): 10 to 12

FLASH POINT: N/A (WATER-BASED LIQUID)

■ PACKAGING

Chlorine Sanitizer is available in 4x1 gallon cases and 5 gallon pails.

■ ADDITIONAL INFORMATION

If you need additional information, please contact our corporate office at 816-471-3602. For a Material Safety Data Sheet go to www.chemicaluniverse.com.

■ DIRECTIONS

Remove all gross food particles/soil from areas to be sanitized with a good detergent, pre-flush, pre-soak or pre-scrape treatment. Rinse with potable water. To sanitize after pre-cleaning, prepare a solution by adding 2 ounce of product to 1 gallon of water. Prepare fresh solution daily or with more frequent use. To sanitize mobile items such as drinking glasses and eating utensils, immerse completely in solution for a least 60 seconds. Remove items, drain from surface and air dry. **POTABLE WATER RINSE IS REQUIRED.**

■ SAFE STORAGE & HANDLING

KEEP OUT OF REACH OF CHILDREN. CORROSIVE. Can cause eye damage and skin burns. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Wear goggles, rubber gloves, and protective clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center immediately for treatment advice. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything to an unconscious person.

IF INHALED: Move person to fresh air. Call a poison control center or doctor for further treatment advice.